

SCHIZOPHRENIA

The WHO estimates that over

21
million

people suffer from schizophrenia, making it one of the top 10 causes of disability worldwide.³



Schizophrenia

is among the most financially costly illnesses in the world.⁴

The disease is marked by positive symptoms

hallucinations
and **delusions**



and negative symptoms

blunted emotions and
social withdrawal⁵

Schizophrenia affects people regardless of race, culture or social class. It typically starts in early adulthood, from age

20

but it can develop at any age from late teens onwards.⁶

The risk of an individual developing schizophrenia during his or her lifetime is approximately.⁶



Schizophrenia is one of Lundbeck's key disease areas. Schizophrenia is a chronic, severe and disabling psychiatric disorder. The disease is characterized by distortions in thinking, perception, emotions, language, sense of self and behaviour. Common experiences include hearing voices and delusions.¹ Schizophrenia affects both men and women, although men tend to develop the condition slightly earlier in life.²

1) <http://www.who.int/mediacentre/factsheets/fs397/en/>

2) www.schizophrenia.com/szfacts.htm.

3) The WHO World Health Report: New understanding, New hope, 2001.

4) Rössler W, Salize HJ, van Os J, Riecher-Rössler A. Size of burden of schizophrenia and psychotic disorders. *Eur Neuropsychopharmacol* 2005; 15 (4): 399-409, Lindström E, Eberhard J, Neovius M, Levander S. Costs of schizophrenia during 5 years. *Acta Psychiatr Scand Suppl* 2007; 116 (435): 33-40.

5) <http://www.nimh.nih.gov/health/publications/schizophrenia-booklet-12-2015/index.shtml>

6) Tsuang MT, Faraone SV. Schizophrenia. Second edition. Oxford University Press Inc., New York: 2005